



# Skin Effect Description in Electromagnetism with a Multiscaled Asymptotic Expansion

Gabriel Caloz, Monique Dauge, Erwan Faou, Victor Péron

## ► To cite this version:

Gabriel Caloz, Monique Dauge, Erwan Faou, Victor Péron. Skin Effect Description in Electromagnetism with a Multiscaled Asymptotic Expansion. Asymptotic methods, mechanics and other applications, Aug 2009, Rennes, France. inria-00528527

**HAL Id: inria-00528527**

**<https://inria.hal.science/inria-00528527>**

Submitted on 22 Oct 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Skin-Effect Description in Electromagnetism with a Scaled Asymptotic Expansion

Gabriel CALOZ

Monique DAUGE

Erwan FAOU

Victor PÉRON

IRMAR

Université de Rennes 1

Macadam Workshop, RENNES

31.08.2009

# References



GABRIEL CALOZ, MONIQUE DAUGE, VICTOR PÉRON (2009)

*Uniform Estimates for Transmission Problems with High Contrast in Heat Conduction and Electromagnetism*




MONIQUE DAUGE, ERWAN FAOU, VICTOR PÉRON (2009)

*Asymptotic Behavior at High Conductivity of Skin Depth in Electromagnetism*

# Transmission Problems with High Contrast Media

## ELECTROMAGNETISM




- $\Omega_-$  Highly conducting material  $\subset\subset \Omega$ : Conductivity  $\sigma_- \equiv \sigma \gg 1$
- $\Sigma = \partial\Omega_-$ : Interface
- $\Omega_+$  Insulating or dielectric body: Conductivity  $\sigma_+ = 0$

Aim: Understand the electromagnetic phenomenon as  $\sigma \rightarrow \infty$

# Transmission Problems with High Contrast Media

## HEAT CONDUCTION



- $\Omega_- \subset\subset \Omega$ : heat conductivity  $a_-$
- $\Sigma = \partial\Omega_-$ : Interface
- $\Omega_+$ : heat conductivity  $a_+$

Aim: Describe the transmission phenomenon as  $|a_-/a_+| \gg 1$

# Outline

**1 Uniform Estimates for Transmission Problems**

**2 3D Multiscaled Asymptotic Expansion**

**3 Numerical Simulations**


# Outline

## 1 Uniform Estimates for Transmission Problems

## 2 3D Multiscaled Asymptotic Expansion

## 3 Numerical Simulations

# Transmission Problems in High Contrast Media



- Heat transfer equation ( $\mathbf{P}_{\underline{a}}$ ):  $\operatorname{div} \underline{a} \operatorname{grad} \varphi = f$

$$\underline{a} = (a_+, a_-) \quad \text{s.t.} \quad |a_-|/|a_+| \rightarrow \infty$$

- Maxwell equations ( $\mathbf{P}_{\underline{\sigma}}$ ):

$$\operatorname{curl} \mathbf{E} - i\omega\mu_0 \mathbf{H} = 0 \quad \text{and} \quad \operatorname{curl} \mathbf{H} + (i\omega\varepsilon_0 - \underline{\sigma}) \mathbf{E} = \mathbf{j}$$

$$\underline{\sigma} = (\sigma_+, \sigma_-) \quad \text{s.t.} \quad \sigma_+ = 0 \quad \text{and} \quad \sigma_- \equiv \sigma \rightarrow \infty$$



# Issues

- ❶ *Uniform piecewise estimates and regularity* in Sobolev norms for solutions  $\varphi$  of  $(\mathbf{P}_{\underline{a}})$
- ❷ *Uniform  $L^2$  estimates* for solutions  $(\mathbf{E}, \mathbf{H})$  of  $(\mathbf{P}_{\underline{\sigma}})$

## Hypothesis

$\Sigma$  is a bounded Lipschitz surface in  $\mathbb{R}^3$

Scalar case:



H.P. HUY, E. SANCHEZ-PALENCIA (1974)

*Limit problem and Strong convergence results as  $a_- / a_+ \rightarrow \infty$*



S. HASSANI, S. NICAISE, A. MAGHNOUJI (2009)

Maxwell case:



H. HADDAR, P. JOLY, H.N. NGUYEN (2008)

*Uniform estimates under a stronger regularity assumption on  $\Sigma$*

# Scalar Problem

## Variationnall Problem

$$V_D = H_0^1(\Omega) \quad \text{or} \quad V_N = \{\varphi \in H^1(\Omega) \mid \int_{\Omega} \varphi \, d\mathbf{x} = 0\}$$

(**VP**<sub>a</sub>): Find  $\varphi \in V = V_D$  or  $V = V_N$  such that

$$\begin{aligned} \forall \psi \in V, \quad \int_{\Omega_+} a_+ \nabla \varphi^+ \cdot \nabla \bar{\psi}^+ \, d\mathbf{x} + \int_{\Omega_-} a_- \nabla \varphi^- \cdot \nabla \bar{\psi}^- \, d\mathbf{x} = \\ - \int_{\Omega} f \bar{\psi} \, d\mathbf{x} + (a_+ - a_-) \int_{\Sigma} g \bar{\psi} \, ds \end{aligned}$$

### Hypothesis (**H**<sub>a</sub>)

- ❶  $f \in L^2(\Omega) \quad \text{and} \quad g \in L^2(\Sigma)$
- ❷  $\int_{\Omega} f \, d\mathbf{x} = \int_{\Sigma} g \, ds = 0 \quad \text{if} \quad V = V_N \quad \text{and} \quad \int_{\Sigma} g \, ds = 0 \quad \text{if} \quad V = V_D$

# Uniform Estimates

## Theorem

Let us assume that  $a_+ \neq 0$ . There exist  $\rho_0 > 0$  independent of  $a_+$  such that for all  $a_- \in \{z \in \mathbb{C} \mid |z| \geq \rho_0 |a_+|\}$ , and for all data  $(f, g)$  satisfying  $(\mathbf{H}_{\underline{a}})$ ,  $(\mathbf{VP}_{\underline{a}})$  has a unique solution  $\varphi \in V$ , which is piecewise  $H^{3/2}$  and

$$\|\varphi^+\|_{\frac{3}{2}, \Omega_+} + \|\varphi^-\|_{\frac{3}{2}, \Omega_-} \leq C_{\rho_0} (|a_+|^{-1} \|f\|_{0, \Omega} + \|g\|_{0, \Sigma})$$

with  $C_{\rho_0} > 0$ , independent of  $a_+$ ,  $a_-$ ,  $f$ , and  $g$ .



GABRIEL CALOZ, MONIQUE DAUGE, V. PÉRON (2009)

Key of the proof: asymptotic expansion for  $\varphi$  with respect to the powers of  $\rho^{-1} := a_+ (a_-)^{-1}$  and convergence of these series in the piecewise  $H^{3/2}$ -norm

# Remarks

- ❶ Similar estimate when the roles of  $a_+$  and  $a_-$  are exchanged. More precise estimate where  $a_+$  and  $a_-$  play symmetric roles
- ❷ The assumption that  $\Sigma$  is Lipschitz is necessary
- ❸ For *polyhedral* Lipschitz interface  $\Sigma$ , uniform piecewise  $H^s$  estimates:

$$\|\varphi^+\|_{s,\Omega_+} + \|\varphi^-\|_{s,\Omega_-} \leq C_{\rho_0} (|a_+|^{-1} \|f\|_{s-2,\Omega} + \|g\|_{s-\frac{3}{2},\Sigma})$$

for  $s \leq s_\Sigma$  with some  $\frac{3}{2} < s_\Sigma \leq 2$

- ❹ For *smooth* interface  $\Sigma$ , uniform piecewise  $H^s$  estimates for any  $s \geq 2$

# Maxwell Problem

## Framework and Boundary Conditions

$$(\mathbf{P}_{\underline{\sigma}}) \quad \operatorname{curl} \mathbf{E} - i\omega\mu_0 \mathbf{H} = 0 \quad \text{and} \quad \operatorname{curl} \mathbf{H} + (i\omega\varepsilon_0 - \underline{\sigma}) \mathbf{E} = \mathbf{j} \quad \underline{\sigma} = (0, \sigma)$$

$$\textcircled{1} \quad \mathbf{E} \times \mathbf{n} = 0 \quad \text{and} \quad \mathbf{H} \cdot \mathbf{n} = 0 \quad \text{on} \quad \partial\Omega$$

$$\textcircled{2} \quad \mathbf{E} \cdot \mathbf{n} = 0 \quad \text{and} \quad \mathbf{H} \times \mathbf{n} = 0 \quad \text{on} \quad \partial\Omega$$

$$\textcircled{1} \quad \text{B. C. 1:} \quad \mathbf{j} \in H(\operatorname{div}, \Omega) = \{\mathbf{u} \in \mathbf{L}^2(\Omega) \mid \operatorname{div} \mathbf{u} \in L^2(\Omega)\}$$

$$\textcircled{2} \quad \text{B. C. 2:}$$

$$\mathbf{j} \in H_0(\operatorname{div}, \Omega) = \{\mathbf{u} \in H(\operatorname{div}, \Omega) \mid \mathbf{u} \cdot \mathbf{n} = 0 \quad \text{on} \quad \partial\Omega\}$$

### Hypothesis (SH)

*The angular frequency  $\omega$  is not an eigenfrequency of the problem*

$$\left\{ \begin{array}{ll} \operatorname{curl} \mathbf{E} - i\omega\mu_0 \mathbf{H} = 0 \quad \text{and} \quad \operatorname{curl} \mathbf{H} + i\omega\varepsilon_0 \mathbf{E} = 0 & \text{in } \Omega_+ \\ \mathbf{E} \times \mathbf{n} = 0 \quad \text{and} \quad \mathbf{H} \cdot \mathbf{n} = 0 & \text{on } \Sigma \\ \text{B.C.1 or B.C.2} & \text{on } \partial\Omega \end{array} \right.$$

# Maxwell Problem

## Uniform $L^2(\Omega)$ Estimate

### Theorem

*Under Hypothesis (SH), there are  $\sigma_0$  and  $C > 0$ , such that for all  $\sigma \geq \sigma_0$ , the Maxwell problem  $(\mathbf{P}_{\underline{\sigma}})$  with B.C.2 and  $\mathbf{j} \in \mathbf{H}_0(\operatorname{div}, \Omega)$  has a unique solution  $(\mathbf{E}, \mathbf{H})$  in  $\mathbf{L}^2(\Omega)^2$ , which satisfies:*

$$\|\mathbf{E}\|_{0,\Omega} + \|\mathbf{H}\|_{0,\Omega} + \sqrt{\sigma} \|\mathbf{E}\|_{0,\Omega_-} \leq C \|\mathbf{j}\|_{\mathbf{H}(\operatorname{div}, \Omega)}$$

*Similar result for B.C.1 and  $\mathbf{j} \in \mathbf{H}(\operatorname{div}, \Omega)$*



GABRIEL CALOZ, MONIQUE DAUGE, V. PÉRON (2009)

Application: Convergence of asymptotic expansion at high conductivity

# Maxwell Problem

## Proof of Maxwell Uniform Estimate

### Lemma

*Under Hypothesis (SH), there are  $\sigma_0$  and  $C_0 > 0$  such that if  $\sigma \geq \sigma_0$  any solution  $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$  of problem  $(\mathbf{P}_{\underline{\sigma}})$  with B.C.2 and data  $\mathbf{j} \in \mathbf{H}_0(\operatorname{div}, \Omega)$  satisfies the estimate*

$$\|\mathbf{E}\|_{0,\Omega} \leq C_0 \|\mathbf{j}\|_{\mathbf{H}(\operatorname{div}, \Omega)}$$

*Similar statement for B.C.1 and  $\mathbf{j} \in \mathbf{H}(\operatorname{div}, \Omega)$ .*

Keys of the proof:

- Electrical Field Decomposition into a regular field and a gradient field



C. AMROUCHE, C. BERNARDI, M. DAUGE, V. GIRAULT (1998)

- Scalar Estimates used for the gradient field  $\nabla\varphi$

# Outline

1 Uniform Estimates for Transmission Problems

**2 3D Multiscaled Asymptotic Expansion**

3 Numerical Simulations



# References and Notations

Asymptotic expansion as  $\sigma \rightarrow \infty$  of solutions of  $(\mathbf{P}_{\underline{\sigma}})$  when  $\Sigma$  is smooth:



E. STEPHAN, R.C. McCAMY (1983-84-85)

*Plane Interface and Eddy Current Approximation*



H. HADDAR, P. JOLY, H.N. NGUYEN (2008)

*Smooth Interface and Impedance boundary conditions*



V. PÉRON (2009)

*Smooth Interface and Perfectly Conducting Electric or Magnetic Wall B.C.*

## Hypothesis

- ①  $\Sigma$  is an oriented smooth compact surface
- ② Perfectly Conducting Magnetic Wall b.c.
- ③  $\mathbf{j} \in \mathbf{H}_0(\text{div}, \Omega)$  and  $\mathbf{j} = 0$  in  $\Omega_-$
- ④ Hypothesis (SH) on  $\omega$

# Asymptotic Expansion

$$\delta := \sqrt{\frac{\omega \varepsilon_0}{\sigma}} \rightarrow 0 \quad \text{as} \quad \sigma \rightarrow \infty$$

By Theorem there exists  $\delta_0$  s.t. for all  $\delta \leq \delta_0$ , there exists a unique solution  $(\mathbf{E}_{(\delta)}, \mathbf{H}_{(\delta)})$  to  $(\mathbf{P}_{\underline{\sigma}})$

Then it is possible to construct series expansions in powers of  $\delta$ :

$$\mathbf{E}_{(\delta)}^+(\mathbf{x}) \approx \sum_{j \geq 0} \delta^j \mathbf{E}_j^+(\mathbf{x}), \quad \mathbf{E}_{(\delta)}^-(\mathbf{x}) \approx \chi(y_3) \sum_{j \geq 0} \delta^j \mathbf{w}_j(y_\beta, \frac{y_3}{\delta})$$

$$\mathbf{E}_j^+ \in H(\text{curl}, \Omega_+) \quad , \quad \mathbf{w}_j \in H(\text{curl}, \Sigma \times \mathbb{R}_+) \quad \text{profiles}$$

$(y_\beta, y_3)$ : “normal coordinates” to  $\Sigma$  in a tubular neighborhood  $\mathcal{U}_-$  of  $\Sigma$  in  $\Omega_-$   
Similar result for the magnetic field.

# Validation of the asymptotic expansion

## Remainders

Aim: proving estimates for remainders  $\mathbf{R}_{m;\delta}$

$$\mathbf{R}_{m;\delta} := \mathbf{E}_{(\delta)} - \sum_{j=0}^m \delta^j \mathbf{E}_j \quad \text{in } \Omega$$

Evaluation of the right hand side when the Maxwell operator is applied to  $\mathbf{R}_{m;\delta}$ :

$$\left\{ \begin{array}{ll} \text{curl curl } \mathbf{R}_{m;\delta}^+ - \kappa^2 \alpha_+ \mathbf{R}_{m;\delta}^+ & = 0 \quad \text{in } \Omega_+ \\ \text{curl curl } \mathbf{R}_{m;\delta}^- - \kappa^2 \alpha_- \mathbf{R}_{m;\delta}^- & = \mathbf{j}_{m;\delta}^- \quad \text{in } \Omega_- \\ [\mathbf{R}_{m;\delta} \times \mathbf{n}]_{\Sigma} & = 0 \quad \text{on } \Sigma \\ [\text{curl } \mathbf{R}_{m;\delta} \times \mathbf{n}]_{\Sigma} & = \mathbf{g}_{m;\delta} \quad \text{on } \Sigma \\ \text{curl } \mathbf{R}_{m;\delta}^+ \times \mathbf{n} & = 0 \quad \text{on } \partial\Omega \end{array} \right.$$

with  $\alpha_+ = 1$  and  $\alpha_- = 1 + i/\delta^2$

# Validation of the asymptotic expansion

## Estimates for Remainders

$$\|\mathbf{j}_{m;\delta}^-\|_{2,\Omega_-} + \|\mathbf{g}_{m;\delta}\|_{\frac{1}{2},\Sigma} + \|\operatorname{curl}_{\Sigma} \mathbf{g}_{m;\delta}\|_{\frac{3}{2},\Sigma} \leq C_m \delta^{m-1}$$

with  $C_m > 0$  independent of  $\delta$ .

### Theorem

*Under Hypothesis in the framework above, for all  $m \in \mathbb{N}$  and  $\delta \in (0, \delta_0]$ , the remainders  $\mathbf{R}_{m;\delta}$  satisfy the optimal estimates*

$$\begin{aligned} \|\mathbf{R}_{m;\delta}^+\|_{0,\Omega_+} + \|\operatorname{curl} \mathbf{R}_{m;\delta}^+\|_{0,\Omega_+} \\ + \delta^{-\frac{1}{2}} \|\mathbf{R}_{m;\delta}^-\|_{0,\Omega_-} + \delta^{\frac{1}{2}} \|\operatorname{curl} \mathbf{R}_{m;\delta}^-\|_{0,\Omega_-} \leq C'_m \delta^{m+1} \end{aligned}$$

# Electrical Profiles

$\mathbf{W}_j =: (\mathcal{W}_{\alpha,j} ; w_j)$  in coordinates  $(y_\beta, Y_3)$  with  $Y_3 = \frac{Y_3}{\delta}$

$$\mathbf{W}_0 = 0$$

$$\mathcal{W}_{\alpha,1} = -j_{\alpha,0}(y_\beta) e^{-\lambda Y_3} \quad \text{and} \quad w_1 = 0$$

$$\mathcal{W}_{\alpha,2} = \left[ \lambda^{-1} \{ b_\alpha^\sigma j_{\sigma,0} - \mathcal{H} j_{\alpha,0} \} - j_{\alpha,1} + Y_3 \{ b_\alpha^\sigma j_{\sigma,0} - \mathcal{H} j_{\alpha,0} \} \right] (y_\beta) e^{-\lambda Y_3}$$


$$w_2 = -\lambda^{-1} D_\alpha j_0^\alpha(y_\beta) e^{-\lambda Y_3}$$

with

$\mathcal{H}$  mean curvature of  $\Sigma$

$$j_{\alpha,k}(y_\beta) := \lambda^{-1} (\text{curl } \mathbf{E}_k^+ \times \mathbf{n})_\alpha(y_\beta, 0) \quad \text{and} \quad \lambda = \kappa e^{-i\pi/4}$$

# Curl Operator and Levi-Civita Tensor



Levi-Civita Tensor  $\epsilon$ :

$$\epsilon^{ijk} = \epsilon_0(i, j, k) / \sqrt{\det(a_{\alpha\beta}(h))}$$

$a_{\alpha\beta}(h)$  : metric on  $\Sigma_h$  and  $\epsilon_0(i, j, k) = \text{sign of } (i, j, k)$

Curl Operator in Normal Parameterization:

$$\begin{cases} (\nabla \times \mathbf{E})^\alpha = \epsilon^{3\beta\alpha} (\partial_3^h E_\beta - \partial_\beta E_3) & \text{on } \Sigma_h \\ (\nabla \times \mathbf{E})^3 = \epsilon^{3\alpha\beta} D_\alpha^h E_\beta & \text{on } \Sigma_h \end{cases}$$

$D_\alpha^h$  : Covariant Derivative on  $\Sigma_h$

# Magnetical Profiles

$\mathbf{V}_j =: (\mathcal{V}_j^\alpha ; v_j)$  in  $(y_\beta, Y_3)$  coordinates

$$\mathbf{V}_0(y_\beta, Y_3) = H_0(y_\beta) e^{-\lambda Y_3},$$

$$\mathcal{V}_1^\alpha(y_\beta, Y_3) = \left[ H_1^\alpha(y_\beta) + Y_3 \left( \mathcal{H} H_0^\alpha + b_\sigma^\alpha H_0^\sigma \right)(y_\beta) \right] e^{-\lambda Y_3},$$

$$v_1(y_\beta, Y_3) = \lambda^{-1} \mathbf{D}_\alpha H_0^\alpha(y_\beta) e^{-\lambda Y_3}$$

with  $H_j^\alpha(y_\beta) := (\mathbf{n} \times \mathbf{H}_j^+) \times \mathbf{n}^\alpha(y_\beta, 0)$

Comparison with



H. HADDAR, P. JOLY, H.N. NGUYEN (2008)

# Skin Effect and Skin Depth

1D model of *Skin Effect* :  $\{z > 0\}$  = half-space conductor body

$$\|\mathbf{E}_\sigma(z)\| = E_0 e^{-\frac{z}{\ell(\sigma)}}, \quad \ell(\sigma) := \sqrt{\frac{2}{\omega \mu_0 \sigma}} \quad \text{Skin Depth}$$

$$\ell(\sigma) = \delta / \operatorname{Re} \lambda.$$

3D model of *Skin Effect*, using profiles  $\mathbf{V}_0, \mathbf{V}_1$ ,

$$\mathbf{V}(y_\alpha, y_3) := \mathbf{V}_0(y_\alpha, \frac{y_3}{\delta}) + \delta \mathbf{V}_1(y_\alpha, \frac{y_3}{\delta})$$

## Definition

Suppose that for all  $y_\alpha$ ,  $\|\mathbf{V}(y_\alpha, 0)\| \neq 0$ . The *Skin Depth* in a conductor body at conductivity  $\sigma$  and  $y_\alpha$  on a Regular Surface  $\Sigma$  is the positive real  $\mathcal{L}(\sigma, y_\alpha)$  defined on  $\Sigma$  taking the smallest value s.t.

$$\|\mathbf{V}(y_\alpha, \mathcal{L}(\sigma, y_\alpha))\| = \frac{1}{e} \|\mathbf{V}(y_\alpha, 0)\|$$



# Asymptotic Behavior for Skin Depth

## Theorem

Let  $\Sigma$  be a Regular Surface with mean curvature  $\mathcal{H}$ . Suppose that  $H_0(y_\alpha) \neq 0$ . The Skin Depth  $\mathcal{L}(\sigma, y_\alpha)$  has the asymptotic expansion

$$\mathcal{L}(\sigma, y_\alpha) = \ell(\sigma) \left( 1 + \mathcal{H}(y_\alpha) \ell(\sigma) + \mathcal{O}(\sigma^{-1}) \right)$$



MONIQUE DAUGE, ERWAN FAOU, V. PÉRON (2009)

Key of the proof:

$$\|\mathbf{v}(y_\alpha, y_3)\|^2 = \left[ \|H_0\|^2 + 2\delta \operatorname{Re}\langle H_0, H_1 \rangle + 2y_3 \mathcal{H} \|H_0\|^2 + \mathcal{O}(\delta, y_3) \right] e^{-2y_3/\ell(\sigma)}$$

# Outline

1 Uniform Estimates for Transmission Problems


2 3D Multiscaled Asymptotic Expansion

**3 Numerical Simulations**

# Numerical Simulations

## FEM Discretization

- Magnetic Field
- Framework:  $\Omega_-$  and  $\Omega$  Axisymmetric Domains
- $\mathbf{F}$  Axisymmetric & Orthoradial  $\rightarrow \mathbf{H}_\sigma$  Axisymmetric & Orthoradial  
2D Problem  $\rightarrow$  accuracy
- Variational Form : Unknown  $H_\sigma = H_{\sigma,\theta}(r, z) \in V$




- Finite Element Library Melina

# Numerical simulations


## Mesh Refinement

$\Omega_-$  &  $\Omega$  coaxial ellipsoidal domains  $\rightarrow \Omega_-^m$  &  $\Omega^m$  coaxial half-ellipsoids




# Numerical Simulations

## Skin Effect in ellipsoidal geometry




$|\operatorname{Im} H_\sigma|$  for  $\sigma = 5 \text{ S.m}^{-1}$




$|\operatorname{Im} H_\sigma|$  for  $\sigma = 80 \text{ S.m}^{-1}$

# Asymptotic Behavior for Skin Depth in Ellipsoidal Geometry

$$|\operatorname{Im} H_\sigma| \text{ for } \sigma = 5\text{S.m}^{-1}$$




$$\text{Conductor} = \Omega_-^m (\mathcal{H} > 0)$$




$$\text{Conductor} = \Omega_+^m (\mathcal{H} < 0)$$

# Asymptotic Behavior for Skin Depth in Ellipsoidal Geometry

$$|\operatorname{Im} H_\sigma| \text{ for } \sigma = 5\text{S.m}^{-1}$$



$$\text{Conductor} = \Omega_-^m$$



$$\text{Conductor} = \Omega_+^m$$


# Numerical Post-Treatment

## Linear Regression


Extract  $|H_\sigma(h)|$  along edges of the mesh of  $\Omega_-^m$  when  $z = 0$ :  $h = 2 - r$ .

Linear Regression in  $skin(\sigma) \rightarrow n(\sigma)$  points  $\rightarrow$  "numerical" slope  $s(\sigma)$

$$\log_{10} |H_\sigma(h)| = s(\sigma)h + b$$



○ :  $\log_{10} |H_\sigma(2 - r)|$  when  $\sigma = 5$   
 $n(\sigma) = 7$ ,  $s(\sigma) = -3,65542$



○ :  $\log_{10} |H_\sigma(2 - r)|$  when  $\sigma = 80$   
 $n(\sigma) = 3$ ,  $s(\sigma) = -16,279162$



# Post-Treatment

## Accuracy of Asymptotics

- Truncated Asymptotic Expansion :  $\mathbf{H}_{\theta,1}^- := \mathcal{H}_{\theta,0}^- + \delta \mathcal{H}_{\theta,1}^-$
- $\log_{10} |\mathbf{H}_{\theta,1}^-(., 2-r)| = c + \beta(\sigma, \Sigma)(2-r) + \mathcal{O}(\frac{1}{\sqrt{\sigma}}; 2-r)$
- "Theoretical" slope  $\beta(\sigma, \Sigma) := \frac{1}{\ln 10} (\mathcal{H} - \frac{1}{\ell(\sigma)})$
- Relative Error between slopes:  $error(\sigma) := \left| \frac{\beta(\sigma, \Sigma) - s(\sigma)}{\beta(\sigma, \Sigma)} \right|$

$\sigma$	5	20	80
$skin(\sigma)$ (cm)	10.3	5.15	2.58
$n(\sigma)$	7	5	3
$s(\sigma)$	-3, 65542	-7, 883903	-16, 279162
$\beta(\sigma, \Sigma)$	-3, 673319	-7, 889506	-16, 32188
$error(\sigma)$ (%)	0, 48	0, 07	0, 26